

THE EMC SPIN CRISIS: COMPARING PROTON AND PHOTON

O.V. TERYAEV

*Joint Institute for Nuclear Research
141980, Dubna, Russia*

February 1, 2008

Abstract

The cancellation of photonic and gluonic anomalies naturally explains the zero singlet contributions to the first moments of proton and photon spin structure functions.

The so-called EMC Spin Crisis, i.e. the small (comparable with zero) value of the singlet axial current proton matrix element, initiated a stream of theoretical papers during the last four years (see, e.g., [1]). I'll discuss, however, only the "anomalous way" of its resolution, exploiting the gluonic triangle anomaly [2]–[4]. Due to its natural interpretation as a large contribution of the polarized gluon density via the anomalous part of the box diagram, it is often thought of [5] as an essentially perturbative QCD approach. The starting point of the analysis, mentioned already in [2] is, however, insensitive to any perfect perturbative approximation. We need only such quite a general result to be valid as the expression of the first moment of the spin dependent structure function g_1^1 through the matrix element of the axial current, the latter being the only gauge-invariant local operator with the required quantum numbers (although either gauge-dependent or nonlocal operators may contribute provided some perfect recipe of small and large distances separation is adopted). On the other hand, it is not conserved due to the axial anomaly. Because of its nonzero anomalous dimension it is a bad candidate for being in the angular momentum sum rule. The required difference is just the anomalous gluon contribution.

Therefore, if the anomalous contribution is present, the large (i.e. not suppressed parametrically) deviation of g_1^1 from the naive expectation is naturally explained. One may ask, however: is it possible to give a natural explanation of the zero result? There are several positive answers [6, 7]. The required cancellation occurs, as a rule, after a more or a less complicated calculation. Here I'll try to show the zero result to be in a direct relationship with an other remarkable phenomenon occurring in the singlet channel, namely, the $U_A(1)$ -problem (the absence of the Goldstone meson in the zero quark mass limit).

Such a relation is already known for a few years [8, 2]: the ghost pole which provides a mass to η' under the mixing with the Goldstone η'_0 generates the additional form

factor $\tilde{G}_2$ of the topological gluon current "nearly forward" matrix element.

$$\langle P, S | K_\mu^5(0) | P + q, S \rangle = 2MS_\mu \tilde{G}_1(q^2) + q_\mu(Sq) \tilde{G}_2(q^2), \quad (1)$$

$$q^2 \tilde{G}_2(q^2)|_0 \neq 0. \quad (2)$$

One should compare this equation with the gauge invariant axial current matrix element case:

$$\langle P, S | J_\mu^5(0) | P + q, S \rangle = 2MS_\mu G_1 + q_\mu(Sq) G_2, \quad (3)$$

$$q^2 G_2|_0 = 0. \quad (4)$$

The conserved quark-gluon current matrix element (its relation to the quark spin is more convenient than the naive one) is expressed via the low-energy η'_0 constants and is comparable with unity.

$$\Delta\Sigma = \frac{\sqrt{N_F}}{2M} f_{\eta'} g_{\eta'_0 NN}. \quad (5)$$

This equation shows that the mentioned profound relation between the Spin Crisis and $U_A(1)$ problem does exist. However, it is (i) again of the numerical origin and (ii) doesn't explain the zero EMC result. Moreover, there is an approach [9], in which the zero q behaviour of G_2 and the occurrence of massless particles are not related. The main argument is the finite zero q limit of $q^\mu(Sq)/q^2$ provided P and $P + q$ are on the same mass shell. My first point is to show that it is not the case: the massless pole is abandoned also by the kinematical arguments.

I start with the following expression for the quark-gluon angular momentum density

$$M^{\mu,\nu\rho} = \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} J_{S\sigma}^5 + x^\nu T^{\mu\rho} - x^\rho T^{\mu\nu}. \quad (6)$$

The first term in the r.h.s. is just the canonical quark spin tensor. Note that the energy-momentum tensor here accumulates also the quark orbital momentum as the *total* gluon angular momentum. We may proceed further along this way and express the quark spin in the orbital form with the simultaneous change of the energy-momentum tensor discovered by Belinfante long ago

$$M^{\mu,\nu\rho} = x^\nu T_B^{\mu\rho} - x^\rho T_B^{\mu\nu}. \quad (7)$$

As the conservation of the angular momentum

$$\partial_\mu M^{\mu,\nu\rho} = 0 \quad (8)$$

immediately leads to the symmetry of $T^{\mu\rho}$, the latter implies that

$$\epsilon_{\mu\nu\rho\alpha} M^{\mu,\nu\rho} = 0. \quad (9)$$

One should conclude that the totally antisymmetric quark spin tensor is somehow cancelled and doesn't contribute to the total angular momentum [10]. Is it nevertheless possible to extract any quantitative information about it?

To do this, one may substitute (6) to (7) and (9) and eliminate T (or, more exactly, its antisymmetric part) from the obtained system of two equations. In the case of classical fields one obtains:

$$(g_{\rho\nu} g_{\alpha\mu} - g_{\rho\mu} g_{\alpha\nu}) \partial^\rho (J_{5S}^\alpha x^\nu) = 0. \quad (10)$$

It is just the neglect of surface terms necessary to apply the Belinfante procedure. Passing to the quantum operators one should integrate (6)–(9) over the whole 3-dimensional space and switch them between nucleon states with the momenta P and $P + q$. As a result, $\frac{\partial}{\partial x^\mu}$ is substituted by $-iq_\mu$ and x^μ by $i\frac{\partial}{\partial q_\mu}$ acting on $\delta(\vec{q})$. The latter is, by definition, equal, up to sign, to the derivative acting on the matrix element. One should obtain straightforward, instead of (10), the following restriction:

$$q^\mu \frac{\partial}{\partial q^\alpha} |_0 \langle P | J_{5S}^\alpha | P + q \rangle = q^\alpha \frac{\partial}{\partial q^\alpha} |_0 \langle P | J_{5S}^\mu | P + q \rangle. \quad (11)$$

To make it more clear, let us multiply both sides by q_μ .

$$q^2 \frac{\partial}{\partial q^\alpha} \langle P | J_{5S}^\alpha | P + q \rangle = (q^\beta \frac{\partial}{\partial q^\beta} - 1) q_\gamma \langle P | J_{5S}^\gamma | P + q \rangle. \quad (12)$$

This equality is obviously valid up to the second and higher powers of q . Note that the differential operator in the r.h.s. clearly subtracts the terms linear in q from the divergence matrix element proportional to sq for the pure kinematical reasons. The G_2 pole term, however, provides a contribution linear in q to the l.h.s. This contradiction is manifested already in (10): the G_2 pole contribution to the r.h.s. is zero, while to the l.h.s. is non-zero.

It is a "kinematical" solution of the $U_A(1)$ problem: the special feature of the singlet channel is its relation to the nucleon spin, rather than existence of the anomaly-induced ghost contribution [11]. The zero mass pole is thus abandoned as both by the kinematical and dynamical analysis, in contradiction with [9].

These deep relations are, however, insufficient to explain why the deviation from the naive value leads to the zero result. One should incorporate the dynamics of axial anomaly. I'll try to show that it is sufficient to suppose the famous t'Hooft consistency principle [12] to be valid: the anomalies should be the same for the fundamental and composite particles.

Let us start with a more clear case of the photon matrix elements. The axial current on-shell matrix element is described by the single form factor

$$\langle \gamma(P, \mu) | J_5^\alpha | \gamma(P + q, \nu) \rangle = G_\gamma(q^2) q^\alpha \epsilon^{\mu\nu\rho\sigma} P_\rho q_\sigma. \quad (13)$$

Axial anomaly manifests itself in the zero quark mass limit as the massless pole in G_γ [13]. At the hadron level the t'Hooft consistency principle requires either massless baryons (to be inserted into the triangle loop) or mesons (manifested in the simplest pole diagram). It is the second possibility which is realized in the Nature. The only exception is the singlet channel: the mentioned $U_A(1)$ problem leads to the conservation of the axial current in the chiral limit. Taking into account gluonic and photonic anomalies one concludes that their contributions should cancel each other. As a result, one obtains the low-energy QCD theorem [14]:

$$\langle \gamma(P) | \frac{\alpha}{2\pi} F \tilde{F} | \gamma(P + q) \rangle + \langle \gamma(P) | \frac{\alpha_s}{2\pi} G \tilde{G} | \gamma(P + q) \rangle = 0. \quad (14)$$

I would like to stress that this equality immediately leads to:

$$\langle \gamma(P, \mu) | J_{5S}^\alpha | \gamma(P, \nu) \rangle = 0. \quad (15)$$

It may be called the *Photon Spin Crisis*. In contrast to the proton one it seems to be very natural from the kinematical point of view. The zero q limit of (13) (note that q^2 is zero only if all its components are zero, provided both photons are on-shell) may be non-zero only due to the anomalous pole. The cancellation of the gluonic and photonic anomalies in the singlet channel just restores the naive result. As to the non-singlet channel the anomalous pole provides the calculable values of the forward axial current matrix element. As a result, one obtains the sum rule for the photon spin structure function g_1^γ [15]:

$$\int_0^1 g_1^\gamma(x) dx = -N_C \frac{\langle e^4 \rangle}{2} \frac{\alpha}{2\pi}. \quad (16)$$

It has been recently criticised [16] for the following two main reasons.

First, for the on-shell photons and massive quarks the r.h.s. of the sum rule is zero. I would like to mention that it was stated already in [15] (see eq.(4) of this Ref.), where the reason for the zero result, namely, the cancellation of the "normal" and anomalous contributions, was identified. It seems that to observe the anomalous contribution the photon virtuality should exceed m_π (not m_ρ , as stated in [16]) that replaces m_q when confinement is taken into account. As it is well known, the m_π is proportional to m_q and turns to zero simultaneously with it, while m_ρ is determined by the condensates $\langle G^2 \rangle$ and $\langle \bar{q}q \rangle^2$ and is independent of the light quark mass. The expected virtuality dependence was discussed in the case of the conformal anomaly contribution to $\gamma\gamma$ -scattering [17]. The anomalous contribution can also be isolated making the transverse momentum cutoff, analogously to the anomalous gluon contribution in the proton case [4]. The latter method seems to be more realistic for the experimental verification of the sum rule.

Second, the specific feature of the singlet channel was missed in [15]. However, the ratio of the meson coupling constants for the singlet and non-singlet channels, appearing in the sum rule r.h.s. [16], may be eliminated. The t'Hooft consistency principle and the cancellation of the gluon and photon anomalies lead to the zero singlet channel contribution which results in the sum rule:

$$\int_0^1 g_1^\gamma(x) dx = -N_C \frac{\langle e^4 \rangle}{2} \frac{\alpha}{2\pi} (1 - c) \quad (17)$$

Here $c = \langle e^2 \rangle^2 / (N_F \langle e^4 \rangle)$ is the ratio of singlet and nonsinglet weights [16]. Note that the zero singlet channel contribution does not imply the zero decay width of $\eta'_0 \rightarrow \gamma\gamma$ because of the ghost pole which substitute the anomalous one.

Passing to the hadron matrix element case one should confront the problem of dealing with *two* above mentioned form factors instead of one. Although the situation here is not yet clear enough, it is possible to give the simple diagrammatic explanation of the EMC result.

The Photon Spin Crisis means that the total transition amplitude of the singlet axial current to two photons is zero provided the direct annihilation $q\bar{q} \rightarrow \gamma\gamma$ (Fig. 1a) and that via a gluon pair (Fig. 1b) are taken into account.

Note that the anomaly in the dispersion [13] approach is generated by the horizontal quark propagator whose "dynamical" smallness ($O(m^{-2})$) for the forward scattering compensates the "kinematical" smallness of the nominator ($O(m^2)$) due to the chirality breaking. The axial current vertex is then just the electron-positron source with the




Figure 1: Cancellation of the photonic (a) and gluonic (b) anomalies of the singlet axial current

required parity. Let us therefore "cut off" [18] the upper part of Fig. 1, passing to Fig.2.

Note that Fig. 2a manifests just the quark contribution to the proton polarized DIS while Fig. 2b the anomalous gluon one (the dashed box represents the hadron-parton transition). The contributions of Fig. 2a and 2b should cancel just like that of Fig. 1a and 1b. It is the diagrammatic explanation of the EMC Proton Spin Crisis.

The photon-gluon blob (Fig. 1b,2b) should be of nonperturbative nature to cancel α_s additional to Fig. 1a,2a. It was studied in the framework of the $1/N_C$ -expansion [19], where the validity of the Leutwyler- Shifman theorem, Fig.1, was checked. Note, however, that this result has a perturbative counterpart. The running coupling constant suppression in the asymptotically free gauge theory means, in practice, the logarithmic decrease in comparison with the Born term, provided the energy scale tends to infinity. This suppression is absent for the anomalous gluon contribution [2], as soon as the Leading Log corrections (i.e. the ladder diagrams) are taken into account. The corresponding anomalous dimension is positive and equal to the one-loop beta-function, compensating the leading order α_s evolution. The partonic interpretation is the α_s^{-1} (nonperturbative-like) growth of the gluon distribution first moment. The reason for this unusual perturbative behaviour is just the Adler-Bardeen non-renormalization theorem. As the nonperturbative effects are also generated by the Axial Anomaly, the coincidence of perturbative and nonperturbative results is just the manifestation of the t'Hooft consistency principle.

The relation of the simplest quark handbag diagram, Fig 2a, to the electromagnetic axial anomaly is probably the most unexpected result. To understand it, let us write down the contribution of Fig. 2a in the following way:

$$T^{\mu\nu} \sim m \epsilon^{\mu\nu\alpha\beta} s_\alpha q_\beta \frac{1}{q^2 + 2(Pq)} = \xi_\parallel \epsilon_\perp^{\mu\nu} \frac{(Pq)}{q^2 + 2(Pq)} = \xi_\parallel \epsilon_\perp^{\mu\nu} \frac{\omega}{2(1-\omega)}. \quad (18)$$

Here m is the mass, s the covariant and $\xi_\parallel$ the longitudinal polarization of the struck quark, $\epsilon_\perp$ is a two-dimensional antisymmetric tensor in the hyperplane orthogonal to




Figure 2: Cancellation of the quark (a) and gluon (b) contributions to the nucleon spin structure function first moment

P and q . Expanding this expression in the unphysical region in powers of ω one notices that the constant term is absent. The same is of course true for the crossed antiquark diagram. Therefore, the Fig 2a contribution to the first moment of g_1 structure function is zero. This fact is usually conspired, because normally the expansion is performed *after* extracting the kinematical factor in the nominator. However, it should be taken into account in the free-field theory: it arises due to the boost in the quark (not hadron!) momentum direction which removes m from the first line in (18).

The anomaly manifests itself if one put q^2 to zero and immediately obtain the ω -independent constant. It cannot be extracted directly in the cross-section because the crucial pole in the quark propagator is changed to δ -function when the discontinuity is taken. The situation is analogous to the QED low-energy theorems case: the amplitude is determined by the pole of the intermediate state propagator and its integral is expressed via the square of the elastic form factor. Therefore we expect that the intensively discussed Gerasimov–Drell–Hearn (GDH) sum rule (see e.g. [20] and ref. therein) in the massless fermion case is closely related to the axial anomaly. The zero value is then expected for the singlet contribution to this sum rule in the close analog with the Photon and Proton Spin Crisis. This hypothesis is strongly supported by the fact that the transition from the Bjorken to GDH sum rule is smooth in the nonsinglet channel [20]: no dramatic sign change is observed. It is interesting that for the second nonsinglet $SU(3)$ combination both Bjorken and GDH sum rules change sign preserving the possibility of smooth interpolation. However, the transition Q_0 is about 4 times lower in this case. It is possibly due to the fact that the additional $\langle mq\bar{q} \rangle$ quark condensates should be taken into account. Whereas the light quark condensate $\langle q\bar{q} \rangle^2$ leads to $G_A/G_V \geq 1$ [21], the light and strange quarks contribution should partially cancel each other to obtain the experimental value of G_A/G_V in this channel. The perturbative result may be continued then to the lower Q region.

There is another possibility to obtain a smooth transition to $Q^2 = 0$, realized recently [22]. If one decompose the GDH sum rule into the contributions of tensor structures $\epsilon^{\mu\nu\alpha\beta} s_{\alpha} q_{\beta}$ and $(sq)\epsilon^{\mu\nu\alpha\beta} P_{\alpha} q_{\beta}$ (known as Schwinger sum rules) the first one

approach zero smoothly. The strong Q^2 dependence is then described by the elastic contribution to the second Schwinger sum rule which is nothing else than the known Burkhardt–Cottingham sum rule for the G_2 structure function. This leads to the zero longitudinal lepton–proton asymmetry at $Q^2 \sim 0.2 GeV^2$.

This approach is qualitatively supported by the fact that the $\Delta(1232)$ saturating about 80% of the GDH integral contribute *only* via the second G_2 tensor if only the dominant magnetic dipole transition form factor G_M is nonzero. The contribution to the first tensor arises only via the interference of the Coulomb quadrupole form factor G_C with the electric quadrupole one G_E and G_M .

One may therefore expect that G_2 is somehow related to the gluon anomaly, as well as the G_M of the $\Delta(1232)$. It is not very strange because both these structures are produced by the helicity-flip, the only source in massless QCD being just the axial anomaly. I would like also to compare the EMC and GDH integrals: both are positive in a chiral limit and are reduced to zero (EMC) and negative value (GDH). The identification of the anomaly in the later case, however, requires further investigation.

I conclude with the statement that the EMC Spin crisis seems to be a natural consequence of the t’Hooft consistency principle realized via the cancellation of the QED and QCD anomalies. The problem, however, requires further investigation, both theoretical and experimental. The latter includes e.g. search for the Photon Spin Crisis, the neutron spin structure investigation and the GDH sum rule Q^2 -dependence.

References

- [1] A.V.Efremov, Preprint CERN-TH.6466/92.
- [2] a) A.V.Efremov, O.V.Teryaev, Preprint JINR E2-88-287.
b) A.V.Efremov, J.Soffer, O.V.Teryaev, Nucl.Phys. **B346**, 97(1990).
- [3] G.Altarelli, G.G.Ross, Phys.Lett. **B214**, 381(1988).
- [4] R.D.Carlitz, J.C.Collins, A.H.Mueller, Phys.Lett. **B214**, 229(1988).
- [5] A.E.Dorokhov, N.I.Kochelev, Yu.A.Zubov, Argonne Preprint PHY-7056-TH-92(1992).
- [6] S.Forte, E.V.Shuryak, Nucl.Phys. **B357**, 153(1991).
- [7] J.H.Kuhn, V.I.Zakharov, Phys.Lett. **B252**, 615(1990).
- [8] A.V.Efremov, J.Soffer, N.Törnqvist, Phys.Rev. **D44**, 1495(1991).
- [9] A.V.Kisselev, V.A.Petrov, Preprint CERN-TH.6355/91.
- [10] R.L.Jaffe, A.Manohar, Nucl.Phys. **B337**, 509(1990).
- [11] D.I.Dyakonov, M.I.Eides, ZhETF **81**, 434 (1981).
- [12] G.t’Hooft in *Recent Development in Gauge Theories*, New York, Plenum Press, 1980.

- [13] a) A.D.Dolgov, V.I.Zakharov, Nucl.Phys.**B27**, 525(1971).
b) A.D.Dolgov, V.I.Zakharov, I.B.Khriplovich, Pis'ma ZhETF **45**, 511(1987).
- [14] H.Leutwyler, M.Shifman, Bern University Preprint BUTP-89/2(1989);
M.A.Shifman, Uspekhi Fiz. Nauk **157**, 561(1989).
- [15] A.V.Efremov, O.V.Teryaev, Phys.Lett. **B240**, 200(1990).
- [16] S.Narison, G.M.Shore, G.Veneziano, Preprint CERN-TH.6467/92.
- [17] A.S.Gorski, B.L.Ioffe, A.Yu.Khodjamirian, Phys.Lett. **B227**, 474(1989).
- [18] O.V.Teryaev, ICTP Preprint IC/91/390(1991), JINR Preprint E2-93-25.
- [19] K.Ghosh, B.Patel, Columbia University Preprint CU-TP-509(1991).
- [20] B.L.Ioffe, V.A.Khoze, L.N.Lipatov, *Hard Processes*, North-Holland, 1984.
J.Soffer, Preprint CPT-90/P.2443(1990).
- [21] V.M.Belyaev, I.I.Kogan, Pis'ma ZhETF **37**, 611(1983).
- [22] J.Soffer, O.Teryaev, Preprint CPT-92/P.2836(1992).